

Development and validation of an instrument to measure informational needs of breast cancer patients in Sri Lanka

D.C. Kuruppu¹, C.N. Wijeyaratne², Nalika Gunawardane³, I. Amarasinghe⁴

¹Medical Library, Faculty of Medicine, University of Colombo, Sri Lanka.

²Professor in Reproductive Medicine, Department of Obstetrics & Gynecology, Faculty of Medicine, University of Colombo, Sri Lanka.

³National Professional Officer (Health Systems Evidence and Analysis), World Health Organisation Country Office, Sri Lanka.

⁴Former Senior Oncological Surgeon, National Cancer Institute, Maharagama, Sri Lanka.

ARTICLE INFO

Article history:

Submission 15.03.2017

Acceptance 15.06.2017

Available online

DOI:

<http://doi.org/10.4038/cjms.v54i1.4821>

Keywords:

Informational needs

Breast cancer

Construct validity

Reliability

Abstract

Background: Breast cancer is the second most common cancer in the world and the commonest in Sri Lanka. The provision of relevant and accurate information in a social-culturally appropriate manner will lead to proper understanding of the disease and treatment options.

Objectives: The aim of this study was to develop and validate the Sri Lankans' Informational Needs Assessment Questionnaire of Breast Cancer (SINAQ-BC).

Methods: Item generation was through review of literature and using qualitative research techniques among stakeholders. Thus developed and pretested 69 items SINAQ-BC was administered through interviews to 150 women at least 18 years of age who have been diagnosed as having breast cancer within the period of one year prior to the study and Exploratory Factor Analysis (EFA) was used for further item reduction. Internal consistency reliability and acceptability were also assessed.

Results: Exploratory factor analysis determined that 58 items should be retained in the questionnaire and that they be best organized into five subscales labelled as disease, diagnosis, treatment, physical care and psychosocial care. SINAQ-BC showed good internal consistency reliabilities with Cronbach's alpha of 0.76 to 0.93. An average 30 minutes (+5 mins) was taken to complete the SINAQ-BC and it did not require any additional explanations demonstrating its acceptability.

Conclusion: The SINAQ-BC demonstrated to be a valid and reliable tool to assess informational needs of breast cancer patients. It is recommended to be used to determine informational needs of women with breast cancer and as a tool to assess effectiveness of educational interventions.

Introduction

Breast cancer (BC) is the second most common cancer in the world and the commonest cancer among women. An estimated 1.67 million new cancer cases were diagnosed in 2012. Incidence rates vary nearly four-fold across the world regions,

with rates ranging from 27 per 100,000 in Middle Africa and Eastern Asia to 92 in Northern America [1]. In Sri Lanka, breast cancer is the commonest cancer among women. The age standardized incidence rate (ASR) of BC in Sri Lanka was 23.0 per 100,000 population in 2010. The crude incidence rate (CR) of the BC has increased by approximately 1 per 100,000 population from the levels in 2009. One in every 40 females in Sri Lanka has a risk of developing BC during their lifetime (0-74 years) [2].

Corresponding Author:

D.C. Kuruppu, Medical Faculty, Faculty of Medicine, Kynsey Road, Colombo 08.

kuruppudc@gmail.com

 <http://orcid.org/0000-0003-0763-0497>



This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Upon diagnosis and during the life thereafter, BC causes a significant emotional impact upon the lives of patients as well as family and friends, work colleagues and the greater community. Many studies reveal that information plays an extremely important role in patients' ability in coping with the disease and their quality of life. Fredette in 1995 [3] stated that information on BC requires the afflicted woman to be well informed to understand her diagnosis, to develop a positive attitude towards the life amidst the threatening implications of the cancer, and to cope well with the condition. Provision of relevant and accurate information in a social-culturally appropriate manner will lead to proper understanding of their treatment options and outlook they need to adopt towards a healthy future [4]. It is therefore important to assess the information needs of women with BC since such an assessment has not been done in Sri Lanka.

Measuring of informational needs require a tool which should be culturally appropriate. A tool to measure informational needs of BC, named as Toronto Informational Needs Questionnaire (TINAQ), has been developed in the European region [5] but none have been developed for Asian Region. Therefore, development of an appropriate instrument to assess informational needs of BC survivors to be used in the Sri Lankan context was considered of paramount importance. It is imperative that validity of the survey instrument to measure informational needs of BC patients be also determined. This study aimed to develop a questionnaire to measure the informational needs of a BC patient and to determine its validity and reliability to assess the needs of information related to diagnosis, treatment modalities and follow up care.

Methods

Development of the tool

The development of the tool in the form of an interviewer administered questionnaire followed a step wise process.

First step was generation of items. Items were identified through review of literature and through focus group discussions. Focus group discussions were carried out with BC survivors, the lay people and the oncologist counselor attached to counselling unit in the National Cancer Institute, Maharagama,

and other staff, to find out suitability of selected items and identify the new items. Sixty nine items were generated from a number of sources including findings in the literature and opinion from the multi-disciplinary panel comprising experts as well as stakeholders i.e. oncologists, community physicians, radiologists, physiotherapists, nutritionist, medical counsellors, social counsellors, nurses and cancer patients.

In the second step, a consultative meeting was organized at which the same panel grouped the items into different subscales or domains that have been identified from literature survey [5]. Using collective professional experience, items that were thought to be repetitions and culturally inappropriate were removed by consensus. Items were then grouped into eight such subscales or domains of informational needs namely, diseases, diagnosis, investigative tests, treatments, physical care, psychological, care family and finance.

The third step was to convert the items into questions and to develop a draft interviewer administered questionnaire to assess breast cancer patients' information needs. The researchers initially drafted the questionnaire in English. The questionnaire was designed to assess the information needs by inquiring the subjects to rate the importance of each item on a five point scale of 1 (not important), 2 (slightly important), 3 (moderately important), 4 (very important) and 5 (extremely important).

Translation of the tool

Two translators, both with a high level of proficiency in English and Sinhala, independently translated the questionnaire into Sinhala. The two Sinhala translated versions were then reviewed by an independent expert proficient in both English and Sinhala languages. The items which were agreed upon were accepted for forward translation. Another two bilingual translators, both with a high level of proficiency in English and Sinhala, independently translated the provisional forward translation back into English without referring to the original English version. The two English translated versions were then reviewed by an independent expert, proficient in both English and Sinhala languages. Same way was used to translate the questionnaire into Tamil using two translators both with high level of proficiency in English and Tamil.

Then questionnaire thus developed was named as Sri Lankans' Information Needs Assessment Questionnaire of Breast Cancer (SINAQ-BC).

Pretesting of the SINAQ-BC

Pretesting the SINAQ-BC was done among ten BC patients at the Cancer Unit of Teaching Hospital in Galle, to assess the understanding and the clarity in the wording of questions. Administration of each questionnaire was followed by a structured interview conducted by the PI. Based on the interviews, a few modifications were done to the instructions stated for some items.

Validation of the SINAQ-BC

The SINAQ-BC together with its instructions was shown to a panel of experts, not involved in developing the study instrument, to assess the content validity. The experts agreed that the content assessed by the SINAQ-BC would be a good reflection to improve patients' knowledge on their disease and services provided for them. The experts also decided that content was subjectively appeared on the SINAQ-BC, confirming its content validity and face validity.

In appraising construct validity, Exploratory Factor Analysis (EFA) using principal component analysis (PCA) was used to assess whether SINAQ-BC had any redundant items and also to assess the suitability of its seven subscales structural. Internal consistency reliability of the SINAQ-BC was assessed using Cronbach's alpha. Bartlett's test of sphericity and Kaiser-Meyer – Olkin measure was performed to assess sampling adequacy.

Study Population

The study population was women at least 18 years of age who have been diagnosed as having BC within the period of one year prior to the study.

Sample Size and sampling technique of the validation study

Nunnally [6] opines that an initial sample of at least 100 subjects is necessary to provide sufficient data for item analysis of a tool being validated. The researcher further says that domain's (subscales) internal consistency, reliability estimates can be made with five subjects per item. The largest subscale in the SINAQ-BC has 19 items with a minimum sample required being 95 (1:5).

Considering these references, 150 subjects were required since this instrument has 69 total items.

Patient visiting outpatient clinics and / or admitted at National Cancer Institute, Maharagama were invited to participate in the study. The patients were not approached on the first day of any new treatment for ethical reasons. Written informed consent was obtained from each patient. A total of 170 BC patients were invited and 156 consented. Interviews were conducted at the convenience of the patient.

Data Collection

The SINAQ-BC developed above was administered by the PI after the interview.

Data Analysis

Exploratory Factor Analysis (EFA) is essential to determine underlying constructs for a set of measured variables. Exploration for redundant items in the SINAQ-BC was based on the results of the exploratory Factor analysis. Following two criteria were used to determine whether an item would be retained.

1. Items with average inter-item correlation of less than 0.20 as the low correlation indicate the item may not be related to topic of interest.
2. When two items had inter-item correlations of 0.80 or higher, indicate that the two items measured the same informational need.

Using the results of EFA, the subscale structure of the SINAQ-BC was also reviewed. Inter correlations of the subscales to total scale and the inter-correlations of the subscales were examined to confirm the suitability of the subscales that were decided upon. Internal consistency reliability for the SINAQ-BC and its subscales was assessed using Chronbach's alpha.

Acceptability was assessed using the average time taken to complete the interview and the response rates. All analysis was carried out using a statistical package for Social Science Version 21.0 (SPSS 21.0) by PI.

The project was approved by the Ethics Committee in the Faculty of Medicine, University of Colombo, Sri Lanka [EC-10-035].

Results

Validation of Sri Lankans' Informational Needs Assessment Questionnaire of Breast Cancer (SINAQ-BC)

The response rate was 96.2%. Age range of the study units were 27 – 70years with mean age of 56.28 years (SD =± 12.87; 95% CI = 54.20 – 58.36). The majority were married (70%) had completed either GCE (O/L) or GCE (A/L) (64%, n=96) and 9% (n= 9) possessed a degree or higher qualification. Approximately half (49%, n=74) reported an average household income of Rs. 20,000 – Rs. 30,000. Only 34% had only undergone surgery as the treatment modality for breast cancer. Thirty three percent had undergone chemotherapy and surgery while balance 33% had undergone radiation therapy and surgery.

Exploratory Factor Analysis

Exploratory Factor Analysis performed by principal component analysis using data. Sampling adequacy was confirmed by Barlett’s test of Sphericity which was significant at $p < 0.01$ (chi square = 1735.4, df = 105, $p = 0.000$ and Kaiser-Meyer-Olkin measure of sampling adequacy which was 0.697, being well above the accepted level of 0.6).

Nine items were deleted because their average inter-item correlations were below 0.20.

Finally the SINAQ-BC contained 58 items. All 58 items had item-total correlations of 0.25 or higher, levels considered acceptable by Nunnally [6]. In all retained items, the means fell above the midpoint of the Likert scale (1-5) with standard deviations in the range of 0.4 – 1.7.

Analyzed 58 items in SINAQ-BC, indicated five subscales. The Eigen values ranged from 1.020 – 5.074. The total variance explained was 68.1%. In the Exploratory Factor Analysis, five subscales were formed out of the seven original subscales in the SINAQ-BC where the two subscales diagnosis and investigative tests were combined as diagnosis and the two subscales on family and finances were combined into a subscale of psychological care. Thus, titles of the subscales after combining, the number of items of each subscale with the relevant factor loadings are shown in Table 1.

Table 1: SINAQ-BC by its subscales with the retained items and the factor loadings

Title of the subscale of information needs	No. of items	items	Factor Loading
		The changes that can be seen on breast	.794
		Detail of the places where they offer breast screening services (whether they are free of charge, such services in my residence, is a referral or an appointment needed)	.758
		What one should do, if a breast abnormality is detected	.758
		Who are at high risk of getting breast cancer	.725
		What causes breast cancer	.716
Disease	14	The age at which the women should get screened for cancer	.669
		How to detect whether I have any other cancer	.663
		Ways of detecting the cancer early	.629
		About breast lumps- why they occur, where they occur, how to recognize	.580
		Whether all women with risk get the cancer	.563
		Information about mammography screening	.538
		Common breast pathologies	.482
		What can women do to minimize their risk of getting breast cancer	.453

		Detail of the places where they offer mammography services (whether they are free of charge, such services in my residence, is a referral or an appointment needed)	.393
		Why each of the confirmatory tests are necessary	.648
		How to prepare for such test	.634
		Whether there are different stages of breast cancer and what do they mean	.590
		Whether the tests can identify the stage of it	.551
Diagnosis	9	How women will act in response of either positive or negative result in each of the above tests.	.545
		The investigations that confirm the diagnosis of breast cancer	.544
		The reasons why doctors suggest certain additional tests. Eg; X – rays, bone scans etc.	.491
		How my illness could affects my life in the future	.480
		what should a breast cancer victim discuss with family / friends about the illness, if diagnosed	.463
		What side effect I should report to the doctor / nurse	.891
		If I have side effects how to deal with them	.887
		If there are ways to prevent side effects / ease treatment.	.828
		The possible side effects of treatment	.824
		The need to be in the hospital for treatment / if so, how long	.805
		How long will I require treatment	.743
		Whether the treatment may give up any time duration.	.703
Treatment	15	About treatment for breast cancer	.695
		Mode of treatment is given	.657
		Whether treatment should do continuously	.647
		How to decide particular treatment plan	.621
		Any food restriction /important food, during treatment	.612
		Any restriction to take other drug (if any), during treatment	.603
		Detail of the places where they offer treatment (whether they are free of charge, such services in my residence, is a referral or an appointment needed)	.543
		Importance of mental preparation for treatment	.505
		When should the breast be examined after surgery	.822
		How to care for my wound or incision	.786
Physical care	5	When should I commence my usual physical activities	.784
		Any additional therapies required to improve recovery after surgery.	.733

		How long will it take to heal	.716
		How to tell if the cancer has come back.	.791
		How to manage mental stresses successfully	.783
		What are the symptoms to get admitted, after treatment?	.772
		How to face the society successfully	.759
		To be aware other family members about their risk for breast cancer	.735
		What are the symptoms to come to the clinic immediately?	.711
Psychosocial care	15	Detail of the places where they offer counselling services (whether they are free of charge, such services in my residence, is a referral or an appointment needed)	.719
		Detail of the places where we are able to buy brassieres & the cost	.698
		Telephone numbers for public (hotline) which can be used to get information on Breast Cancer	.638
		How often should be followed up in respect to treatment mode?	.631
		How should family members discuss about the disease with the patient at an initial level of diagnosis	.533
		Information on financial support	.467
		Where my family can go if they need help to deal with my illness	.451
		Importance of continue treatment & care	.441
		Behaviour of newly diagnosed women	.422

As shown in the Table 1 all the retained items contained a factor loading of 0.393 or more.

Table 2 shows that the inter correlations of the subscale to total scale ranged from 0.77 – 0.89 and

the inter-correlations of the SINAQ-BC subscale scores ranged from 0.47– 0.84. This confirmed that each subscales best correlated with the total tool compared to its correlation to other subscales.

Table 2: Inter-correlation of SINAQ-BC total and subscale scores (n=150)

Subscale	Total	Disease	Investigation	Treatment	Physical care	Psychosocial care
Disease	.85	-				
Investigation	.88	.74	-			
Treatment	.89	.84	.77	-		
Physical care	.77	.47	.55	.60	-	
Psychosocial care	.89	.58	.71	.64	.80	-

Internal consistency reliability

Results of internal consistency reliability of the SINAQ-BC are shown in Table 3. Cronbach's

alpha for the total and each of the five subscales indicated values 0.76 – 0.95 indicating the high reliability.

Table 3: Internal consistency reliability of the SINAQ-BC

	Cronbach's Alpha
Total scale	0.95
Subscales	
Disease	0.93
Diagnosis	0.78
Treatment	0.86
Physical	0.76
Psychosocial	0.85

Acceptability of SINAQ-BC

None of the women had difficulty in responding to the SINAQ-BC and the meantime taken to complete the interview was 30 minutes (± 5 mins) and all were able to respond to the items without additional explanations.

Discussion

The SINAQ-BC is the only validated instrument developed to assess informational needs of BC patients in Sri Lankan context. It is expected to function as a valid measure to identify information needs which facilitate designing of the patient educational programmes.

The existing literature is not rich enough to provide a sound conceptual foundation for informational needs of BC patients. As Flick(1998) and Neuman (1997) have suggested that focus group is useful in exploratory research or in generating new ideas to develop hypotheses [7,8]. Hence, the present study used both these methods to generate items of the SINAQ-BC.

A hospital-based cross sectional study design was employed to assess the validity and reliability of the SINAQ-BC. Thus, purposive sampling technique was considered as the best technique to ensure representation of women with an initial diagnosis of breast cancer in the treatment phases of surgery, chemotherapy, radiation therapy.

To be considered as acceptable items in the questionnaire, item analysis was completed as done by Nunnally [6]. Items with item-total correlations of 0.25 or higher levels was considered as an acceptable item in the questionnaire, correlation below 0.20 considered as not interesting items and if two items have inter-item correlation of 0.80 or higher, it considered as same informational need, hence, retained only one item. Rarely both items were retained if they seem to be dealing with different informational needs. Content validity of the instrument was established as emphasized by Latwin in 1994 to assess the content completeness of the instrument [9].

Nunnally [6] opines that an initial sample of at least 100 subjects is necessary to provide sufficient data for item analysis to decide whether the research should be continued or abandoned. The researcher further says that domain's (subscales) internal consistency, reliability estimates can be made with five subjects per item. Accordingly, the largest subscale in the SINAQ-BC has 19 items, a minimum sample would be 95 (1:5). Ferguson & Cox [10] stated at their pre-analysis that minimum participants to variable ratio would be in between 2:1 – 6:1. Considering these references, 150 subjects were used for the study since this instrument has 69 total items.

Factor analysis is a complex statistical procedure, which essentially reduces a large set of variables into much smaller set of variables with common characteristics [11]. As such, it is an empirical

method of determining 'the underlying dimensionality' of a set of variables [12]. Factor analysis suggested by Galloway et al. (1997) [5] for the Toronto Informational Needs Questionnaire-Breast Cancer (TINQ-BC) was used in the present research.

Reliability refers to the dependability and repeatability of an instrument. Wilkin et al. [13] explained that if an instrument is more reliable, there is lower element of random error. Reliability also provides an estimate of homogeneity. Cronbach's α is used to estimate the internal consistency reliability of SINAQ-BC. The alpha coefficient for each of the subscales and the total scale was above 0.80, which is considered as a satisfactory estimate of reliability by Burns & Grove [14]. Nunnally also said that this is the level necessary for instruments used in basic research [6]. A tool (TINQ-BC) developed by Galloway et al in 1997 [5] also showed that overall alpha coefficient was 0.97 while the corresponding figure for SINAQ-BC was 0.95. Domain α of TINQ-BC ranged from 0.73 to 0.93 while for SINAQ-BC it was 0.76 to 0.93.

Conclusion

The SINAQ-BC demonstrated to be a valid and reliable tool to assess informational needs of BC patients in relation to diagnosis, treatment modalities and follow up care. This is evident by high internal consistency, reliability, content validity and construct validity. Its validity to assess informational needs was confirmed irrespective of the treatment modality that the patient was undergone. It is recommended that the SINAQ-BC be used to determine informational needs of women with BC in Sri Lanka and as a tool to assess educational interventions to improve knowledge and coping of BC among patients.

Acknowledgements

Acknowledge University of Colombo for funding of the study [Research grant AP/3/2012/PG/05]. The authors are thankful to the Director of National Cancer Institute, Maharagama, Director of Teaching Hospital in Galle and all health professionals in these two hospitals for the immense support rendered.

References

1. Estimated cancer incidence, Mortality and Prevalence Worldwide in 2012. GLOBOCAN 2012: http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx (accessed 28 Feb 2017).
2. Cancer incidence data: Sri Lanka 2010. 12th Publication. Colombo: National Cancer Control programme, 2016.
3. Fredette, S.L. Breast cancer survivors: concerns and coping. *Cancer Nurse*. 1995; 18: 35-46.
4. Johnson JD, Meischke H. Women's preferences for cancer information from specific communication channels. *American Behavioural scientist*. 1991; 34: 742-755.
5. Galloway S., Graydon J., Harrison D., et al. Informational needs of women with a recent diagnosis of breast cancer: development and initial testing of a tool. *Journal of Advanced Nursing*. 1997; 25:1175-1183.
6. Nunnally J.C. *Psychometric Theory*; 2nd edition, McGraw-Hill Book Company, Toronto, 1978: 278-282.
7. Flick, U. *An introduction to qualitative research*. Sage: London, 1998.
8. Neuman, W.L. *Social research methods – qualitative and quantitative approaches*, 3rd ed., Allyn and Bacon, Boston, MA, 1997.
9. Litwin MS. Measuring health related quality of life in men with prostate cancer. *Journal of Urology*. 1994;152:1882-1887.
10. Ferguson, E. and Cox, T. Exploratory Factor Analysis: A Users' Guide. *International Journal of Selection and Assessment*, 1993; 1: 84-94. doi:10.1111/j.1468-2389.1993.tb00092.x
11. Templeton H.R.M. & Coates V.E. Adaptation of an instrument to measure the informational needs of men with prostate cancer. *Journal of Advanced Nursing*, 2001; 35(3), 357-364.
12. Polit D.F. *Data Analysis and Statistics for Nursing Research*, Prentice Hall, London, 1996.
13. Wilkin D., Hallam L. & Doggett A.M. *Measures of Need and outcome for Primary Health Care*, Oxford University Press, Oxford.1992.

14. Burns N. & Grove S.K. *Understanding Nursing Research*, W.B. Saunders Co. : London, 1995.